



## **Uptake and depuration of three differently functionalized zinc oxide nanoparticles to *Daphnia magna***

**Skjolding, Lars Michael; Winther-Nielsen, Margrete; Baun, Anders**

*Publication date:*  
2012

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Skjolding, L. M., Winther-Nielsen, M., & Baun, A. (2012). *Uptake and depuration of three differently functionalized zinc oxide nanoparticles to Daphnia magna*. Poster session presented at 7th International Conference on the Environmental Effects of Nanoparticles and Nanomaterials, Alberta, Canada.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Uptake and depuration of three differently functionalized zinc oxide nanoparticles to *Daphnia magna***

Lars Michael Skjolding<sup>1</sup>, Margrethe Winther-Nielsen<sup>2</sup>, Anders Baun<sup>1</sup>

<sup>1</sup>Department of Environmental Engineering, Technical University of Denmark, Building 113, Kgs. Lyngby, Denmark

<sup>2</sup>DHI, Agern Allé 5, DK-2970 Hørsholm, Denmark

**Abstract:**

During the last couple of years the use of nanoparticles (NP) has dramatically increased. Zinc oxide nanoparticles (ZnO NP) have a wide range of applications e.g. in personal care products, paints and semi conductors. However, few studies have so far investigated the ecotoxicity of ZnO NP and to our knowledge the bioaccumulation behavior in regards to difference in functionalization of ZnO NP has not been studied previously. In this study, experiments with the crustacean *Daphnia magna* was carried out to test if the changes in lipophilicity of functionalized ZnO NP would respectively increase or decrease the bioaccumulation compared to ZnO NP. The experiments were based on ISO 6341 *Daphnia* immobilization test and bioaccumulation studies with *Daphnia magna* including a 24h uptake and 24h depuration phase. The tested ZnO NP was of same primary size (35 nm) but with different functionalizations (ZnO, ZnO-OH, ZnO-C<sub>8</sub>H<sub>17</sub>). Characterization included ICP-MS, DLS, BET and TEM. Preliminary results show a fast uptake of ZnO NP ( $k_1=0.11\text{ h}^{-1}$ ) and fast depuration ( $k_2=-0.23$ ). In contrast, ZnO-OH did not show significant uptake compared to control. The poster will focus on trends in uptake and depuration of NP with changes of functionalization in terms of lipophilicity compared to non-functionalized ZnO.